

Agronomic evaluation of Tunisian accessions of chili pepper (*Capsicum frutescens* L.)

Kaouther ZHANI¹, Wissem HAMDY¹, Sami SEDRAOUI², Rami FENDRI¹, Oula LAJIMI¹, Chérif HANNACHI¹

¹ Higher Institute of Agronomy, 4042, Chott Mariem, Tunisia, Sousse University,

² Faculty of Sciences of Bizerte, Tunisia, Carthage University

Abstract - Five local accessions of chili pepper (*Capsicum frutescens* L.): *Tébourba*, *Somâa*, *Korba*, *Awled Haffouz* and *Souk Jedid*, were cultivated in the experimental station of Higher Institute of Agronomy, Chott Mariem, Sousse (Tunisia) for their agronomic evaluation. Results showed that the pepper accessions were significantly different ($p < 0.05$). *Korba* accession had the superior vegetative characters (stem height, stem width, leaf area, plant height, number of bifurcations) and produced the highest number of flowers (356), fruits (211) and yield (22 t/ha). Fruits of *Tébourba* accession are the longest (118 mm), the widest (15 mm), the heaviest (12 g) and have the thickest (2 mm) pericarp, the longest (96 mm) placenta and the biggest number of seeds (127 seeds).

Key Words: *Capsicum frutescens*, fruits, genotypes, quality, yield.

1. Introduction

Pepper (*Solanaceae* family and *Capsicum* genus) is an important agricultural crop. It is an annual plant grown for its berries extensively used as spice and in medicine. Its fruits, used as spice, vegetable and herbal remedy, are known by their highest vitamin C content among all vegetables [1] and their antioxidant characteristics [2]. It plays an important role in the economy of many countries. It is the world's second important vegetable, ranking after tomatoes and it is the most produced type of spice flavouring and colouring for food while providing essential vitamins and minerals. In 2013, Food and Agriculture Organization (FAO) statistics estimated world production of *Capsicum* peppers at 21.3 million tonnes from a harvested area of 1.6 million ha (i.e. an average yield of 13.4 t/ha) [3]. Comparatively, yield in the developing countries is about 10 – 30% of that in developed countries [4]. However, Tunisia is known to be one of the major producers of pepper in the world.

In fact, in Tunisia, pepper is widely grown in all regions both on open air and under greenhouse and occupies the fourth largest area planted by gardening. In

2014, the area allocated to this crop reached 21,200 ha with a national yield of 18.1 t/ha [5]. Pepper is very appreciated by Tunisian consumer's especially pungent varieties which are used fresh to prepare different dishes, dried (spice) or pickled.

Pepper production in Tunisia is not unconnected with the climatic condition and day length while the moisture content of the soil and the prevailing temperature has important effects on the growth and yield of the crop. In general, the maximum growth and production of pepper occur between a temperature range of 18°C and 30°C [4], in a loam or silt loam soil with good water-holding capacity (but also on many soil types as long as the soil is well drained) and the soil pH between 5.5 and 6.8 [6]. The majority of cultivated varieties are hybrids belonging to the genus *Capsicum annuum* and estimated to be 34, all are officially allowed by Agriculture Minister [71]. The varieties belonging to *Capsicum frutescens* or Chili pepper are local varieties, named "Baarbid". The most production of this species in Tunisia is localized in the region of Korba. Nevertheless, this type of pepper is cultivated in other regions and no study was done about the production of the accessions cultivated there.

Therefore, for the first time, the present research studies the agronomic performance of local Tunisian chili pepper. Thus, five accessions of chili pepper (*Capsicum frutescens* L.) are cultivated under greenhouse in Chott Meriem (Sousse, Tunisia) in order to select the best performing accession(s).

2. Materials and methods

2.1. Experimental protocol

Research was conducted in the experimental station of Higher Institute of Agriculture of Chott Mariem (Tunisia). Five local accessions of chili pepper were included in this study. They were collected from three Tunisian regions: Manouba, Cap Bon and Sidi Bouzid. Two of them came from two locations in Cap Bon; Somâa (Sm) and Korba (Kb), two were from the state of Sidi Bouzid:

Awled Haffouz (Az) and Souk Jedid (Sj) and the fifth was collected from Tébourba (Tb) in Manouba.

The work lasted from January to October 2011. In January 5th seeds of the five accessions were sown in alveolar containers consisting of 104 cells and containing fertilized peat (N-P-K, 12-24-12). These containers were covered by white plastic to accelerate seeds germination and emergence and put under greenhouse until the plants were ready for transplanting. Watering was done based on climatic condition with a fine watering can and hand weeded. Fungicide (Proplant: 200 ml/hl) was applied before the fungal devastation as preventive activity. Before transplanting, farmyard manure (7 t/ha) was dispersed into the soil and land preparation was done mechanically.

At five leaves stage, plants were transplanted on March 3rd under monotunnel greenhouse covered with polyethylene low density plastic. The plants are arranged, randomly with 5 repetitions, in a single line at a density of two plants/m². They were vertically dressed, irrigated regularly using system drip irrigation and fertilized according to the crop needs. Fungicides were used to protect plants against aphids (Talastar: 80 ml/hl), moths and leaf miners (Lannate: 200 ml/hl) and powdery mildew (Solfo: 500 g/hl).

2.2. Data collection

A number of 18 agronomic traits (Table 1) are used to study of the growth and the production of the plant. 60 plants, 30 leaves and 30 fruits within each accession were used to characterize the five genotypes.

2.3. Soil analysis

Soil samples were collected annually from the 0 to 20 cm depth. Soil pH was determined potentiometrically in 1:2 soil/distilled water suspensions after shaking. Soil organic matter was measured using the walkley and black method. The Kjeldahl method was used to determine the soil nitrogen (N). Phosphorus (P) was determined using Olsen's method. Potassium (K) was determined using a flame photometer. Calcium and Mg were determined using atomic absorption spectrophotometer

2.4. Data Analysis

SPSS software 13.00" was used to analyze the data and Duncan's multiple range test was used to separate the means at 5% (p<0.05).

Table 1. Agronomic characters in study

Characters studied
Plant Height (PH) (at red ripening): measured in cm from soil level to the top most growth point of above ground plant
Stem Height (SH)
Time of beginning of flowering (BF): number of days from plantation to the appearance of the first flower on second flowering nods.
Time of maturity : number of days from plantation to apparition of the first mature fruit)
Leaf area (LA)(cm²): determined by planimeter (Metter 3100)
Stem width (SW): measured in cm at widest point
Number of bifurcations (NB) : counted 120 days after plantation
Number of flowers per plant (NF): determinate 120 days after transplantation
Number of fruits per plant (Nbfr)
Yield per plant (Yp) and per hectare (Yh)
Fruit weight (WF) (g) : determinate on SHIMAZDOU A×200 digital scales
Fruit Length (LF): measured in mm with Digital Caliper
Fruit Diameter (DF) : measured in mm with Digital Caliper
Thickness of flesh (Tf): measured in mm with Digital Caliper
Stalk length (Sl): measured in mm with Digital Caliper
Placenta length (Pl) (mm)
Number of seeds per fruit (SN)
100-seed-wight (100 sw) in g

2. Results

Initial soil physicochemical characteristic were show in the Table 2. The value of pH is 7.8 indicated that is a weakly alkaline soil. Soil analyses gave values of 2.7% organic matter (OM), and nutrients 27 (N), 412 (P), 816 (K), 193 (Ca) and 22 (Mg) (mg.kg⁻¹ soil) and it have a sandy loam texture soil.

Table 2. Soil proprieties

Chemical proprieties (mgkg ⁻¹)						
pH	OM (%)	N	P	K	Mg	Ca
7.8	2.7	27	412	816	22	193
Physical proprieties (%)						
Clay	Sand	loam	Field capacity	Wilting point		
23	62	15	7.5-14.2	3.2-8.6		

Examination of agronomic performance in the vegetative stage (Table 3) showed a significant difference ($p < 0.05$) in stem diameter (10.36 to 16.6 cm), leaf area (46.81 to 69.68 cm²) and plant height (56.16 to 114.83 cm). Kb cv demonstrated the best values while Sj cv had the lowest ones. The number of bifurcations, counted 120 DAP, was measured only in Kb, Az, Sm and Tb cv and it was ranging between 168.12 (Az cv) and 228.3 (Kb cv) bifurcations.

Significant variability in flowering time (DAP to the appearance of the first flower) and the total number of flowers per plant was observed (Table 3) and the range was respectively 44 to 58 DAP and 212.7 to 356.5 flowers. Tébourba cv was the earliest to flowering with 44 days while Sm cv took the longest days (58 days). Kb cv had the highest number of flowers per plant (356.5) whereas Tb cv had the lowest number (212.7 flowers).

Table 3. Vegetative parameters of five Tunisian chili pepper accessions

	Tb	Sm	Kb	Az	Sj
SH (cm)	12.30 b	12.73 b	12.89 b	13.74 b	41.40 a
SW (cm)	14.30 b	14.70 b	16.60 a	13.90 b	10.36 c
LA (cm ²)	62.09 b	52.82 c	69.70 a	70.47 a	46.81 d
PH (cm)	102.96 b	104.73 b	114.83 a	90.60 c	56.16 d
NB	207.90 b	194.50 c	228.30 a	168.12 d	-
BF	44.00 b	58.00 a	48.00 b	52.00 b	54.00 a
NF	212.69 d	308.48 b	356.50 a	285.80 c	268.61 c

(a-c): Means in the same line followed by the same letter are not significantly different at 5% level according to Duncan test.

The difference in yield between the five accessions was significant; it is demonstrated in Table 4. The highest fruit yield per plant was obtained from Kb cv with 210.95 fruits while Tb cv had the least fruit yield with 101.28 fruits. The estimation of yield in g per plant showed that Kb cv was the most performing accession (870.61 g) while Sj cv produce the lowest yield per plant (406.8 g). Thus, the yield per hectare ranged from 10.16 (Sj cv) to 21.76 t (Kb cv). Therefore, Kb cv is the most performing accession and it is probably due to its best responses in vegetative growth in terms of height of plant, diameter stem, leaf area, number of bifurcation, fruits number, flowers number. The positive correlation study between yield and vegetative and productive parameters (Table 5) confirm this.

Table 4. Productivity of five Tunisian chili pepper accessions

	Tb	Sm	Kb	Az	Sj
NbFr	101.28 e	195.24 b	210.95 a	168.12 c	149.23 d
Yp (g)	600.4 c	758.5 b	870.61 a	617.26 c	406.78 d
Yh (t)	15.01 c	18.96 b	21.76 a	15.43 c	10.16 d

(a-e): Means in the same line followed by the same letter are not significantly different at 5% level according to Duncan test.

Fruit characteristics showed also a significant difference among the pepper accessions with respect to all the parameters measured (Table 6). Fruits of Tb cv had the highest length (117.9 mm), weight (11.72 g) and placenta length (95.89 mm). Az cv produced the largest (14.67 mm) fruits with the thickest pericarp (2.14 mm). Kb cv had the pedicel with the biggest length (40.23 mm). Sj cv recorded the lowest values in all traits studied. Finally, the number and weight of seeds are studied and a significant difference was moreover recorded. Tb cv produced the higher number of seeds (126.8) while Sj cv produced the lowest one (63.26). Again Tb cv had the highest weight of seeds (0.570 g) whereas Sj cv had the lowest weight (0.432).

Correlation between fruit parameters were also studied (Table 7) and higher positive correlation was observed between weight and length fruit ($r=0,814$), between weight and diameter fruit ($r=0,739$), between placenta length and fruit length ($r=0,571$) and between pericarp thickness and diameter of fruit ($r=0,511$)

Table 5. Correlation coefficients among growth parameters in Tunisian hot pepper

	PH	SH	SW	LA	NB	NF	NbFr	Yp
PH	1							
SH	0,021*	1						
SW	0,094*	0,077	1					
LA	0,052	0,186	0,222*	1				
NB	0,16**	-0,033	0,065	-0,043	1			
NF	0,45**	0,59**	0,150	0,140	0,25**	1		
NbFr	0,547*	0,084	0,267*	-0,184	0,174*	0,052*	1	
Yp	0,246**	0,535*	0,097	0,115	0,099**	0,431*	0,47**	1

Table 7. Correlation coefficients among production parameters in Tunisian hot pepper

	WF	DF	LF	PI	Tf	SN
WF	1					
DF	0,739**	1				
LF	0,814**	0,70**	1			
PI	-0,621**	-0,44**	0,565**	1		
Tf	-0,594**	0,511**	-0,43**	0,81**	1	
SN	-0,671**	-0,59**	-0,62**	0,57**	0,568**	1

Table 6. Fruit characteristics of five Tunisian chili pepper accessions

	Tb	Sm	Kb	Az	Sj
LF (mm)	117.9a	98.38c	82.95d	102.7b	53.51e
DF (mm)	14.67a	12.01b	11.68c	15.63a	8.90d
WF (g)	11.72a	6.04cd	5.57d	10.18b	2.36e
Tf(mm)	2.15ab	1.39c	1.36c	2.42a	1.14d
PI(mm)	95.89a	72.95c	51.97d	87.66b	39.74e
SI (mm)	38.18b	37.48c	40.23a	38.78b	35.38d
SN	126.8a	75.23d	95.53c	106.73b	63.26e
sw (g)	0.533b	0.521b	0.452c	0.570a	0.432d

(a-e) Means in the same line followed by the same letter are not significantly different at 5% level according to Duncan test.

3. Discussion

The review of the agronomic performance of the studied accessions showed a significant difference in the stem height, collar diameter, leaf area and plant height. These results agrees with the findings of many others works on pepper [8,7,10]. The determination of leaf area is an indicator of the power plant since it is directly related to the photosynthetic capacity and chlorophyll synthesis [11], while the diameter of the stem reflects the plant vigor [12] because larger is the diameter better is the plant in supporting best vegetative part and then the fruit load. The height of plant is an important growth character directly linked with the productive potential of plant. An optimum plant height is claimed to be positively correlated with productivity of plant [13,14].

According to the literature, the difference in height of the plant is mainly attributed to the genetic potential [15] but also to environmental factors, especially temperature which must be in order of 25°C to ensure good vegetative growth to plant pepper[16]. Also, The capacity of genotype to absorb soil nutrients from the soil [17] could be a cause in this difference in plant height,

especially nitrogen [18]) and phosphorus [19] which have enhancing effect on the vegetative growth of plants by increasing cell division and elongation.

Our result indicates also significant variations in number of bifurcations. This trait was also used to separate accessions of *Capsicum* and similar variability was observed confirming our results [20]. Generally, this difference is related to a genetic make-up of the crop [21] and/or influence of organic and inorganic nutrients in the soil [17]. The growth temperature can also control the formation of branches as it was confirmed in 10 Tunisian varieties of pepper [22].

Similarly, there was a variation in flowering time and total number of flowers. Analogous result was obtained on 10 local ecotypes in Tunisia [20]. In fact, earliness in flowering is an important parameter to explore because it

The significant difference in yield between the five accessions found in this study agrees with the results of other researchers on pepper [28,29] and other crops like tomato [30], rapeseed [31] and white[32]. The recorded variations of varieties in marketable yield could be due to their differences in genetic make-up and/or agro ecological adaptations compared to the locations in which they had evaluated [33].

With regards to relationship between vegetative growth and yield and yield components of hot pepper, researchers confirmed our results and proved that there is a positive and highly significant correlation between yield and number of branches [34], yield and plant height [35], yield and height and diameter of stem[36] and yield and fruit number [37,38]. High temperatures strongly influence also reproductive development of chili pepper [39] because elevated temperature during post-pollinisation reduced fruit number of pepper [40].

In this study, it was also observed that Tunisian chili peppers accessions are significantly different in their sizes. Similar significant heterogeneity between pepper was discovered in fruits with regards to length and weight [41], diameter[21], pericarp thickness [42], placenta length[20] and pedicel length[43], number of seeds [44].

In our study, a positive correlation has been found between fruit weight and length, and between fruit weight and diameter [42]. Similarly, pericarp thickness is positively correlated with the diameter [34] and weight [46] of fruit. In fact, the fruit weight variability is mostly genetic linked to chromosome 2 [47]. However, it was confirmed[48] that the weight of pepper fruit is controlled by several loci, some with a large effect (locus 2.2 *fw* on chromosome 2) and others with a small effect (locus *fw4* on chromosome 4). Also, temperature [49], plant growth[50], plant nutrition [51], agricultural practices

[52] and competition between fruits that are developing simultaneously on the plant [53] may also affect the growth and the size of pepper fruits.

For seeds, the difference in their number and weight is related to the cultivar [54] on the first hand, to temperature [55] on second hand and to the positive correlation with fruit size on the third hand [56] especially weight [57] and length[35]. In literature, the choice of these characters in the selection of fruits in pepper is related to many reasons. Fruits with thickest pericarp are the most withstand to transportation shocks and have the highest dry matter) [36]. Larger pepper fruits are the most attractive by consumer [58] and so get premium prices. Also, higher is the size of fruits greater is the capability for storage of photosynthates and lower is the rate transpiration [59].

4. Conclusion

Thus, from this study, agronomic variations observed between the five Tunisian *Capsicum frutescens* accessions allow us to identify Korba (Kb) accession as the most performing in production and Tébourba (Tb) as the accession with highest quality of fruits. Then these accessions could be good parents in breeding for improvement programs of pepper.

REFERENCES

- [1] J.A. Osuna-Garcia, M.M. Wall, and C.A. Waddell, "Endogenous levels of tocopherols and ascorbic acid during fruit ripening of New Mexican-type chile (*Capsicum annuum* L.)". *J.Agric.Food Chem.*, vol.46, No.12, pp. 5093-5096, 1998.
- [2] N.Y. Shotorbani, R. Jamei, and R. Heidari, "Antioxidant activities of two sweet pepper *Capsicum annuum* L. varieties phenolic extracts and the effects of thermal treatment". *Avicenna Journal of Phytomedicine*, pp.1-10, 2012.
- [3] FAO, 2015. <http://faostat3.fao.org>
- [4] G.J.H. Grubben, and I. M. Tahir, *Capsicum species*, In: G. J. H. Grubben, and O. A., Denton, (Ed). *Plant Resources of Tropical Africa 2, Vegetables* PROTA Foundation, Wageningen, Netherlands, Backhugs Publishers, Leiden, Netherlands/ICTA, Wageningen, Netherland, 2004.
- [5] CNEDA, Economic budget of the Ministry of Agriculture. National Center for Agricultural Research and Development, Ministry of Agriculture, Tunisia, 2014.
- [6] T. Berke, L.L. Black, N.S. Talekar, J.F. Wang, P. Gniffke, S.K. Green, T.C. Wang, and R. Morris, *Suggested Cultural Practices for Chili Pepper*, AVRDC Publication (Number: 05-620), 2005.
- [7] H.S. Gill, B.M. Asawa, and P.C. Thakur, "Correlation, pathcoefficient and multiple regression analysis in

- sweet pepper". *Indian J Agric Sci.*, vol. 47, No.8, pp. 408-410, 1977.
- [8] M.D. Aljebjo, and F.C. Orakwue, "Characteristics of some pepper populations commonly grown in Nigeria". *Capsicum and Eggplant Newsletter.*, vol. 21, pp. 22-24, 2002.
- [9] A.J. Perez-Lopez, J.M. Lopez-Nicolas, E. Nunez-Delicado, F.M. Del Amor, and A.A. Carbonell-Barrachina, "Effects of agricultural practices on color, carotenoids composition and mineral contents of sweet peppers, cv. Almuden", *J.Agric.Food Chem.*, vol.55, pp. 8158-8164, 2007.
- [10] E.R. Rego, M.M. Rego, F.L. Finger, C.D. Cruz, and V.W.D. Casali, "A diallel study of yield components and fruit quality in chilli pepper (*Capsicum baccatum*)". *Euphytica.*, vol.168, pp. 275-287, 2009.
- [11] J.L. Prioul, and N. Schwebel-Dugu, "Source-sink manipulations and carbohydrate metabolism in maize". *Crop.Sci.*, vol.32, pp. 751-756, 1992.
- [12] H. Sbai, Caractérisation morphologique de quelques écotypes locaux de piment de saison (*Capsicum annum* L.). Master's thesis in Agriculture Durable. Higher Institute of Agronomy, chott Mariem; University of Sousse, Tunisia, 2010.
- [13] L. Aliyu, M.K. Ahmed, and S.G. Ado, "Relationships between some characteristics in chili pepper (*Capsicum frutescens*)". *Capsicum Newsletter.*, vol. 10, pp.47-48, 1991.
- [14] I.N. Saeed, K. Abbasi, and M. Kazim, "Response of maize (*Zea mays*) to nitrogen and phosphorus fertilization under agro-climatic condition of Rawalokot Azad Jammu and Kashmir". *Pak. J. Biol. Sci.*, vol. 4, pp.53-55, 2001.
- [15] B. Aloni, E. Pressman, and L. Karrni, "The effect of fruit load, defoliation and night temperature on the morphology of pepper flowers and fruit shape". *Annals of Botany.*, vol.83, pp.529-534, 1999.
- [16] P. Erard, A. Bellamy, D. Berry, A. Buffière, A. Cappy, J. Dumolin, A. Guillon, C. Hutin, and D. Iazard, *The pepper*. CTIFL (Edt), Paris, 2002.
- [17] W.A. El-Tohamy, A.A. Ghoname, and S.D. Abou-Hussein, "Improvement of pepper growth and productivity in sandy soil by different fertilization treatments under protected cultivation". *J.Applied.Sci.Res.*, vol.2, pp.8-12, 2006.
- [18] R.K.A. Egharevba, and R.K.E. Law-Ogbomo, "Comparative effects of two nitrogen sources on the growth and yield of roselle (*Hibiscus sabdariffa*) in the rainforest region. A case study of Benin City, edo state, Nigeria". *J.Agron.*, vol. 6, pp.142-146, 2007.
- [19] J.G.M. Vos, and H.D. Frinking, "Nitrogen fertilization as a component of integrated pest managements of hot pepper (*Capsicum* sp.) under topical lowland conditions", *Int.J.Pest.Manage.*, vol. 43, pp.1-10, 1997.
- [20] S. Ben Mansour-Gueddes, N. Tarchoun, J.M. Teixeira da Silva, and S. Saguem, "Agronomic and chemical evaluation of seven hot pepper (*Capsicum annum* L.) populations grown in an open field. Fruit", *Vegetable and Cereal Science and Biotechnology.*, vol. 4, pp. 93-97, 2010.
- [21] A. Rivera-Martinez, L. Terren Poves, J.M. Rodriguez Bao, J.L. Andrés Ares and J. Fernandez Paz, "Characterization of local pepper lines from Northwest Spain". *Capsicum and Eggplant Newsletter*, vol. 23, pp. 25-28, 2004
- [22] N. Tarchoun, "Behavioral study of seven varieties and three pepper lines (*Capsicum annum* L.) grown in cold greenhouse", *Tropicultura*, vol.19, N°.1, pp.37-42, 2001.
- [23] H. Demirsoy, and L. Demirsoy, "Characteristics of some local sweet cherry cultivars from homeland", *J.Agron.*, vol.3, pp. 88-89, 2004.
- [24] M. Hassan, M. Ilyas, M. Wasiullah, and M. Khan, "Performance of wheat varieties in Barani areas of Kohat division", *Pak.J.Biol.Sci.*, vol. 4, pp. 114-, 2001.
- [25] P.L. Polowick and V.K. Sawhney, "Temperature effects on male fertility and flower and fruit development in *Capsicum annum* L.", *Scientia Horticulturæ*, vol.25, pp.117-127, 1985.
- [26] K. Alizadeh, and J. Carapetian, "Genetic variation in a safflower germplasm grown in rainfed cold drylands", *J.Agron.*, vol.5, pp.50-52, 2006.
- [27] I. Sreelanthakumary, and L. Rajamony, "Variability, heritability and genetic advance in chilli (*Capsicum annum* L.)", *Journal of Tropical Agriculture*, vol. 42, pp. 35-37, 2004.
- [28] F. Marame, L. Desalegne, H. Singh, C. Fininsa, and R. Sigvald, "Genetic components and heritability of yield and yield related traits in hot pepper", *Res.J.Agric.Biol.Sci.*, vol. 4, pp.803-809, 2008.
- [29] K.E. Law-Ogbomo, and J.E. Law-Ogbomo, "Characterization and evaluation of some cultivars of sweet pepper (*Capsicum annum*)", *Not Sci Biol.*, vol.2, N°. 1, pp. 49-54, 2010.
- [30] T. Pradeepkumar, D.B.M Joy, N.V. Radhakrishnan, and K.C. Aipe, "Genetic variation in tomato for yield and resistance to bacterial wilt", *Journal of Tropical Agriculture*, vol.39, pp. 157-158, 2001.
- [31] P.R. Khoshnazar, M.R. Ahmadi, and M.R. Ghanadha, "A study of adaptation and yield capacity of rapeseed (*Brassica napusum* L.) cultivars and lines", *Iran.J.Agric.Sci.*, vol.31, pp. 341-352, 2000.
- [32] M. Hassan Wasiullah, I. Javed, and I. Mohammad, "Evaluation of wheat varieties under the agro-climatic conditions of barani agricultural research station Kohat", *J.Agron.*, vol.2, pp.8-12, 2003.
- [33] M. Fekadu, and G. Dandena, "Status of vegetable crops in Ethiopia", *Ugandan Journal of Agriculture*, vol.12, n°.2, pp. 26-30, 2006
- [34] D. Seleshi, Evaluation of Elite Hot Pepper Varieties (*Capsicum* species) for growth, dry pod yield and quality under Jimma condition, South West Ethiopia. Master's thesis of Science in Horticulture (Vegetable Science). School of Graduate Studies, College of

- Agriculture and Veterinary Medicine: Jimma University, 2011.
- [35] Z. Yayeh, and A.C. Zeven, "Variation in Yugoslavian hot pepper (*Capsicum annuum* L.) accessions", *Euphytica*, vol.97, pp.81-89, 1997.
- [36] E.R. Rego, M.M. Rego, C.D. Cruz, F.L. Finger, and V.W.D. Casali, "Phenotypic diversity, correlation and importance of variables for fruit quality and yield traits in Brazilian peppers (*Capsicum baccatum*)", *Genet Resour Crop*, vol.58, n°.6, pp. 909-918, 2011.
- [37] H.S. Gill, B.M. Asawa, and P.C. Thakur, "Correlation, pathcoefficient and multiple regression analysis in sweet pepper", *Indian J Agric Sci*, vol.47, n°.8, pp. 408-410, 1977.
- [38] C.R. Gupta, and R.D.S Yadav, "Genetic variability and path analysis in chilli (*Capsicum annuum* Linn.)", *Genetica Agraria*, vol.38, pp. 425-432, 1984.
- [39] P. Pagamas, and E. Nawata, "Effect of high temperature during the seed development on quality and chemical composition of chili pepper seeds", *Japanese journal of tropical agriculture*, vol.51, pp.22-29, 2007.
- [40] E.R. Erickson, and A.F. Makhart, "Flower developmental stage and organ sensitivity of bell pepper (*Capsicum annuum* L.) to elevated temperature", *Plant Cell and Environment*, vol.25, pp.123-130, 2002.
- [41] S. Sood, N. Kumar, K.S. Chandel, and P. Sharma, "Determination of genetic variation for morphological and yield traits in bell pepper", *The Indian Journal of Agricultural Sciences*, vol.81, n°7, pp.590-594, 2011.
- [42] M.M. Moreno, C. Moreno, J. Villena, and I. Mancebo, "Agro-morphological characterization of 16 traditional pepper (*Capsicum annuum* L.) cultivars from Castilla-La Mancha (central Spain). III International Symposium on Plant Genetic, 2010.
- [43] G.O. Nkansah, A. Ayarna, and T.J. Gbokie, "Morphological and yield evaluation of some *Capsicum* lines in two agro-ecological zones of Ghana", *Journal of Agronomy*, vol.10, pp.84-91, 2011.
- [44] G.U. Rao, A. Ben Chaim, Y. Borovsky, and I. Paran, "Mapping of yield-related QTLs in pepper in an interspecific cross of *Capsicum annuum* and *C. frutescens*", *Theor.Appl.Genet*, vol.106, pp.1457-1466, 2003.
- [45] C. Shifriss, J. Zacks, and A. Goldman, "Some notes on the *Capsicum annuum* L." *Euphytica*, vol.46, pp. 275-277, 1989.
- [46] S.D. Lannes, F.L. Finger, A.R. Schuelter, and V.W.D. Casali, "Growth and quality of Brazilian accessions of *Capsicum chinense* fruits", *Scientia Horticulturae*, vol.112, pp.266-270, 2007.
- [47] S. Zygiar, A. Ben Chaim, A. Efrati, G. Kaluzky, and Y. Borovsky, "QTLs mapping for fruit size and shape in chromosome 2 and 4 in pepper and comparison of the pepper QTL map with that of tomato" *Theoretical and Applied Genetics*, vol. 111, pp.437-445, 2005.
- [48] I. Paran, and E. Van der Knapp, "Genetic and molecular regulation of fruit and plant domestication traits in tomato and pepper", *Journal of Experimental Botany*, vol.58, pp. 3841-3852, 2007.
- [49] K. Kato, "Flowering and fertility of forced green pepper at lower temperature", *Journal of the Japanese Society for Horticultural Science*, vol. 58, pp. 113-121, 1989.
- [50] E. Jovicich, J.D. Cantliffe, and J.P. Stofella, "Fruit yield and quality of greenhouse grown bell pepper as influenced by density, container and trellis system", *Hort. Technology*, vol.14, pp.507-513, 2004.
- [51] F.M. Del Amor, "Yield and fruit quality response of sweet pepper to organic and mineral fertilization", *Renewable Agriculture and Food Systems*, vol. 22, pp. 233-238, 2007.
- [52] M. Pérez-Grajolas, V.A. Gonzales-Hernandez, M.C. Mendoza-Castillo, and C. Pena-Valdivia, "Physiological characterization of Manazo hot pepper (*Capsicum pubescens* R, P) landres", *JAMER.SOC.HORT.SCI*, vol.129, n° .1, pp.88-92, 2004.
- [53] A.M. Ali, and W.C. Kelly, "The effects of inter-fruit competition on the size of sweet pepper (*Capsicum annuum* L.) fruits", *Sci. Hort*, vol.52, pp. 69-76, 1999.
- [54] V.S. Kulkarni, Studies on variability, correlation, morphological and molecular diversity in dry chillies (*Capsicum annuum* L.), Master's thesis of Science (Agriculture) in genetics and plant breeding. College of Agriculture, Dharwad, University of Agricultural Sciences, Dharwad. 2006.
- [55] P. Pagamas, and E. Nawata, "Sensitive stages of fruit and seed development of chili pepper (*Capsicum annuum* L.var Shishito) exposed to high temperature stress", *Scientia horticulturae*, vol.117, pp. 21-25, 2008.
- [56] M. Aleemullah, A.M. Haigh, and P. Holford, "Anthesis, anther dehiscence, pistil receptivity and fruit development in the Longum group of *Capsicum annuum*", *Australian Journal of Experimental Agriculture*, vol. 40, pp.755-762, 2000.
- [57] V.M. Russo, "Planting date and plant density affect fruit seed of Jalapeno peppers." *J. Hortsci.*, vol.38, pp. 520-523, 2003.
- [58] T. Beyene, and P. David, "Ensuring Small Scale Producers in Ethiopia to Achieve Sustainable and Fair Access to Pepper Market", *Uganda Journal of Agriculture* vol. 3, n°.2, pp.113-119, 2007.
- [59] J.A. Barrera, M.S. Hernandez, L.M. Melgarejo, O. Martinez, J.P. Fernandez-Trujillo, "Physiological behavior and quality traits during fruit growth and ripening of four Amazonic hot pepper accessions", *J.Sci.Food.Agric*, vol.88, pp.847-857. , 2008.